



09/21/00

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE
REQUEST FOR FILING APPLICATION UNDER 37 CFR 1.53(b)
WITHOUT FILING FEE OR EXECUTED INVENTOR'S DECLARATION**

Assistant Commissioner for Patents
Washington, DC 20231

Atty. Dkt. 121-112

Date: September 21, 2000

Sir:

This is a request for filing a new PATENT APPLICATION under Rule 53(b) entitled:
SURFACE MODIFIED PARTICULATE COMPOSITIONS OF BIOLOGICALLY ACTIVE SUBSTANCES
without a filing fee and/or without an executed inventor's oath/declaration.
This application is made by the below identified inventor(s). Attached hereto are the following papers:

- ☒ An abstract together with
36 pages of specification and claims including
15 numbered claims and also attached is/are
5 sheets of accompanying drawings.

- ☒ This application is based on the following prior provisional application(s):

Application No. **Filing Date**
60/154,964 September 21, 1999

respectively, the entire content of which is hereby incorporated by reference in this application, and priority is hereby claimed therefrom.

- ☒ Please amend the specification by inserting before the first line: --This application claims the benefit of U.S. Provisional Application No. 60/154,964, filed September 21, 1999, the entire content of which is hereby incorporated by reference in this application.--

- | | | | | | |
|----|----------------------|--|--------------------------------|--------|---------------|
| 1. | Inventor: | Gary | W | Pace | Australia |
| | | (first) | MI | (last) | (citizenship) |
| | Residence: (city) | Raleigh | (state/country) North Carolina | | |
| | Post Office Address: | 6913 Hunters Way, Raleigh, North Carolina | | | |
| | (Zip Code) | 27615 | | | |
| 2. | Inventor: | Awadhes | K. | Mishra | India |
| | | (first) | MI | (last) | (citizenship) |
| | Residence: (city) | Montreal | (state/country) Quebec, Canada | | |
| | Post Office Address: | 670 de Gaspe App. 201 Nuns Island Verdun, Montreal, Quebec, Canada | | | |
| | (Zip Code) | H3E 1H8 | | | |

NOTE: FOR ADDITIONAL INVENTORS, check box ☒ and attach sheet with same information.

Address all future communications to NIXON & VANDERHYE P.C., 1100 North Glebe Road, 8th Floor, Arlington, Virginia 22201.

1100 North Glebe Road, 8th Floor
Arlington, Virginia 22201-4714
Telephone: (703) 816-4000
Facsimile: (703) 816-4100
ARC:pfk

NIXON & VANDERHYE P.C.

By Atty: Arthur R. Crawford, Reg. No. 25,327

Signature: 

Reg No 36663

jc905 U.S. PTO
09/667328
09/21/00



3. Inventor: Robert A. Snow Canada
(first) (last) (citizenship)
Residence: (city) West Chester (state/country) Pennsylvania
Post Office Address: 118 Cratin Lane, West Chester, Pennsylvania
(Zip Code) 19380

U.S. PATENT APPLICATION

Inventor(s): Gary W Pace
Awadhesh K. Mishra
Robert A. Snow

Invention: SURFACE MODIFIED PARTICULATE COMPOSITIONS OF
BIOLOGICALLY ACTIVE SUBSTANCES

***NIXON & VANDERHYE P.C.
ATTORNEYS AT LAW
1100 NORTH GLEBE ROAD
8TH FLOOR
ARLINGTON, VIRGINIA 22201-4714
(703) 816-4000
Facsimile (703) 816-4100***

SPECIFICATION

BACKGROUND OF THE INVENTION

Oral, parenteral, and pulmonary delivery of biologically active substances, particularly water-insoluble pharmaceutically active substances, has been addressed by formulating these substances as micron and sub-micron sized particles suspended in water or ultimately present as a suspension in an aqueous environment. Various methods and compositions of such formulations have been reported in literature.

For instance, US Patents 5,091,188; 5,091,187 (D. H. Haynes, "Phospholipid-coated Microcrystals: Injectable Formulations of Water-Insoluble Drugs." Feb. 25, 1992) describe a MicroCrystal technology of delivering undiluted or highly concentrated drug substances as aqueous suspensions of micron- or submicron-sized particles stabilized with physiologically safe, tissue compatible, and pharmaceutically acceptable surface modifiers such as natural and synthetic bipolar lipids. Water-insoluble drugs are rendered injectable by formulation as aqueous suspensions of phospholipid-coated microcrystals. The crystalline drug is reduced to 50 nanometer to 10 micrometer dimensions by sonication or other processes inducing high shear in the presence of phospholipid or other membrane-forming amphipathic lipid. The membrane-forming lipid stabilizes the microcrystal by both hydrophobic and hydrophilic interactions, coating and enveloping it and thus protecting it from coalescence, and rendering the drug substance in solid form less irritating to tissue. Additional protection against coalescence is obtained by a secondary coating by additional membrane-forming lipid in vesicular form associated with and surrounding but not enveloping the lipid-encapsulated drug particles. Tissue-compatible formulations containing drug in concentrations up to 40% (w/v) are described. The preparations can be injected intra-lesionally and in numerous other sites, including intra-venous, intra-arterial, intra-muscular, intra-dermal, etc. The disclosure describes examples of formulations and pharmacokinetic data with antibiotics, anthelmintic drugs, anti-inflammatory drugs, local and general anesthetics, and biologicals. These compositions can be dried and reconstituted into aqueous suspension.

US Patent 5,145,684 describes aqueous suspensions of sub-micron size particles of poorly water-soluble therapeutic or diagnostic agent onto which are adsorbed a non-crosslinked surface modifier.

US patent 5,922,355 discloses aqueous suspensions of surface-modified submicron size particles of pharmaceutical or other water-insoluble or poorly water-soluble substances prepared using a combination of one or more surfactants such as poloxamers, poloxamines,

polyoxyethylene sorbitan fatty acid esters and the like together with natural or synthetic phospholipids. The combination of phospholipids and surfactants allows the formation and stabilization of the sub-micron and micron size surface-modified particles in the form of their aqueous suspensions via hydrophilic, lipophilic and electrostatic interactions and therefore prevent these particles from aggregation or flocculation or particle size growth on storage. These aqueous suspensions of water-insoluble or poorly soluble compounds are manufactured in presence of the surface-modifying agents by a particle size reduction process such as sonication, homogenization, milling, microfluidization and precipitation, or recrystallization and precipitation of the compound using antisolvent and solvent precipitation.

The international publication WO 030615A1 teaches the use of careful selection of surface modifying agents on the basis of their hydrophile-lipophile balance (HLB) to prepare sub-micron and micron-size stable particles of water-insoluble or poorly soluble drugs or other industrially useful insoluble compounds suspended in an aqueous medium containing at least one surface modifier. The surface modifier or modifiers are selected by an HLB-based selection criteria for selecting the type and amount of surface modifiers used to obtain surface-modified sub-micron size stable aqueous suspensions of water insoluble or poorly water-soluble solid drugs.

One of the disadvantages of these technologies is that the surface modified particles of water-insoluble drug substances in aqueous suspension may tend to grow over time because of the dissolution and recrystallization phenomenon known as "Ostwald ripening".

Another disadvantage of these technologies is the requirement of removing water from the surface-modified particle suspensions prior to converting them to suitable solid dosage forms. It is known in the art of aqueous suspensions of surface-modified particles that in many instances water removal processes such as freeze drying and spray drying may cause excessive growth of the particle size or irreversible agglomeration and thereby adversely affect the advantageous properties of the microparticulate composition.

In addition, there are be applications where an aqueous medium is not suitable for surface-modified sub-micron particle suspensions. Alternative medium for such particle delivery systems can be a suitable oil or a suitable hydrophobic liquid or a suitable amphipathic medium in which the substance is substantially insoluble.

Various inventions have been disclosed relating to the suspension of surface modified particles of water-insoluble drugs in non-aqueous media or dispersed as an aqueous suspension within non-aqueous media.

For instance US Patents 5,091,188 and 5,091,187 claim that phospholipid surface modified particles prepared in aqueous suspension and then dried can be dispersed in a pharmacologically acceptable, water-miscible polar organic liquid with a dielectric constant greater than 30, which polar organic liquid does not substantially dissolve the lipid membrane or membranes or the drug substance or a mixture of the polar organic liquid with water. Additionally, a water-immiscible oil can be utilized to facilitate contact between the drug crystals or particles and the primary layer of amphipathic membrane-forming lipid, or to slow the rate of drug dissolution or to otherwise modify the rate of drug release. Also, within their aqueous suspension, the drug microparticles can be precoated by a layer of a waxy phospholipid-compatible solid having a melting point between 37 degree C and 100 degree C before or during the application of the primary layer of membrane-forming amphipathic lipid, the waxy phospholipid-compatible solid selected from paraffin, tristearin, ethyl oleate, cetostearyl alcohol, cetyl alcohol, myristyl alcohol, stearyl alcohol and petrolatum.

US patent 6,086,376 teaches the production of dry suspension aerosol formulation having a density in the range of from 1.0 to 1.5 g/ml consisting essentially of surface stabilized particles of water-insoluble drugs with a mean size range of 0.1 to 10 microns. The surface modifiers consist of one or more membrane-forming phospholipids and at least one surfactant surrounding the drug core. The surface modified particles are dispersed in 1,1,1,2-tetrafluoroethane HFA134a or 1,1,1,2,3,3,3-heptafluoropropane HFA 227 propellant, wherein the density of the coated drug microparticles substantially matches the density of the propellant, and the amount of phospholipid coating on the drug microparticles is more than 0.1% and less than 200% of the weight of the drug. Densities of various components of this invention are carefully controlled to provide a suspension of the surface modified drug microcrystals that upon agitation remains homogeneous for sufficiently long time to allow delivery of the said aerosol formulation. Even though water-free compositions of surface-modified drug microcrystals are suspended in pressurized fluorocarbon propellant, these aerosolizable compositions are neither intended to undergo self-dispersion upon exposing to aqueous environment, nor is there any requirement of exposing these pressurized propellant

based compositions to aqueous medium prior to or during the intended drug delivery mode, *i.e.*, inhalation of the aerosol.

Further, in US patents 5,560,931 and 5,571,536 is described formulations of water insoluble drug compounds as surface modified particles suspended in the aqueous phase of water-in-oil emulsions using digestible oils or fatty acids as the non-aqueous oil phase. The particles are formed by media milling in the presence of the surface modifier. The water (containing surface modified particles)-in-oil emulsion is then formed using a high shear emulsification procedure. The drug must be both water-insoluble and oil-insoluble. Drug insolubility in oil is required; otherwise it will partition into the oil phase by migrating from the water pool to the continuous oil phase and thus will render the formulation out of the scope of invention. This delivery system also suffers from the problem of presence of water that may not allow the use of many popular packaging systems such as hard- or soft-gelatin capsules or starch capsules, because these packaging systems are known to dissolve in aqueous media or their stability is severely compromised by even trace quantities of water. In addition, this technique may be suitable only for high potency drugs because of availability of a very small quantity of water in the oil-in-water emulsion which is the final medium of drug-delivery in this case.

In another method US patent 5,858,398 describes the formation of the particles of both water-soluble and insoluble drugs in the presence of a surfactant and suspending the particles in a micelle form in an oily substance. The microparticles described in this invention contain:

- (1) at least one pharmaceutically-active agent, and
- (2) at least one water soluble or miscible phospholipid, and
- (3) at least one lipid soluble or miscible phospholipid, and
- (4) at least one non-ionic surfactant having an HLB value of about 15 or greater, and
- (5) at least one non-ionic surfactant having an HLB value of about or less, and
- (6) at least one water soluble or miscible sterol compound.

The microparticles of the above composition are suspended in at least one fatty acid having a chain length of C14 or less. The composition may optionally contain at least one fatty acid having a chain length of C16 or greater in a concentration of about 5 w/v % or less.

The composition is prepared by admixing the pharmaceutically-active agent, phospholipids, surfactants, and sterol, micronizing the admixture to form microparticles, and suspending the microparticles in at least one fatty acid of chain length of C14 or less to form microparticles in a micelle. This invention is not designed to specifically address the problems associated with the formulation of water-insoluble drugs, especially stability and potential drug particle growth on exposure to water.

US patent 4,622,219 and US patent 4, 725, 442 describe a micron or sub-micron sized particulate composition of water-insoluble oily drug that is dispersed as an oil-in-water emulsion stabilized by emulsifiers such as natural or semi-synthetic phospholipids.

In another approach to the delivery of water-insoluble drugs, the drug may be dissolved in biocompatible oils. Further these solutions of the drug in oil may be dispersed in water using surfactants to form oil in water emulsions. A high-energy process such as a high-shear emulsification procedure is used to produce oil-in-water emulsions. These compositions require for their formation typically more than 70% of water. For example, US patent 5,660,858 describes pharmaceutical compositions consisting essentially of an oil-in-water emulsion containing a synthetic medium chain triglyceride in which is dissolved a therapeutically effective amount of a cyclosporin, phospholipid, and optionally free fatty acid or a salt thereof, non-ionic surfactant, ionic surfactant, glycerol, salts, buffers, preservative, osmotic modifier and antioxidant.

A special embodiment of oil-in-water type emulsions as carriers of drug substances is so-called solid lipid nanoparticles (SLN) which are prepared by using oils that melt at a temperatures higher than ambient temperature. The liquid oil-in-water emulsion is first made at a suitable temperature and then is cooled down to allow solidification of the oily core of the emulsion droplet. See Siekmann *et al.*, Pharm. Pharmacol Lett., **1**, 123-126 (1992) and Muller *et al.*, European patent no. 0,605,497 (1996).

Oil-in-water emulsions are limited by the following:

- (i) Solubility of the pharmaceutically or biologically active substance in oil or hydrophobic agent may not be enough that a solution can be prepared for making an oil-in-water emulsion using a preferably small quantity of oil or other hydrophobic agents.

- (ii) A large volume of the dispersion (either oil-in-water or water-in-oil types) is necessary for delivering the required amount of the biologically active substance. Such quantities of dispersions cannot be packaged in convenient manner, for example, in hard- or soft-gelatin capsules, or in other types of capsules, or in coated tablets or pills.
- (iii) Further, presence of water in these dispersions may not allow the use of many packaging systems such as hard- or soft-gelatin capsules or starch capsules because these packaging systems are known to dissolve in aqueous media.
- (iv) In many cases the oil-in-water emulsions in the form of solid lipid nanoparticles are known to suffer from partial expulsion of the hydrophobic drug during a cooling step of their preparation which adversely affects the entrapment efficiency of the carrier system.

The formulation of both water-soluble and insoluble drugs in oil in water emulsions where the drug is dispersible or soluble in the oil has been applied to achieve sustained release of the drug.

In this regard the publication WO 041682A1 describes a lipophilic microparticle having an average particle size ranging from 0.1 to 200 micrometers, comprising a lipophilic substance and an active ingredient selected from the group consisting of a protein or peptide drug and an antigen, retains the full activity of the active ingredient, and when formulated in the form of an oil dispersion or oil-in-water emulsion, it releases in an in vivo environment the active ingredient in a controlled manner over a long period.

In another approach to the formulation of water-insoluble drugs the drug may be dissolved in water miscible organic solvents.

For example, US patent 6,046,163 describes therapeutic compositions comprising from 0.1 to 20% by weight of a water-insoluble cyclosporin and a vehicle comprising from 1 to 60% by weight of polyethers constituted by, preferably, substances like the water miscible ethoxy diglycol or polyethylene glycols 300 to 600 and/or substances like preferably dimethyl isosorbide, dimethyl isodide and dimethyl isomannide, and from 1 to 60% by weight of a mixture of glyceryl monoesters of C8-C22 fatty acids and hexaglyceryl to pentadecaglyceryl monoesters of C8-C22 fatty acids in a ratio of 1:2 to 1:6.

In another approach the water-insoluble drug is dissolved in a mixture of oils and surfactants such that on addition to water the components spontaneously emulsify or disperse

to form small particles of the oily phase in which the drug is dissolved. These small oil-phase particles are known as emulsion or microemulsion droplets. Microemulsions, also known as self-dispersing systems, differ from emulsions (which are heterogeneous dispersions of a discontinuous phase of droplets in a continuous phase of bulk liquid) by being as "a system of water, oil, and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution" (see Danielsson and Lindman, "The definition of microemulsion" *Colloids Surf.* 3: 391 (1981)). Oil-in-water type microemulsions are known to form spontaneously, without addition of energy, from microemulsion preconcentrates by mixing with aqueous component. Microemulsion (or emulsion) preconcentrates are compositions of the entire oil-in-water type microemulsion (or emulsion) components except water packaged in a convenient dosage form such as a dilutable and/or ingestible solution or capsule or gelcap.

In US patent 5,932,243 Fricker *et al.* teach about a pharmaceutical composition containing macrolide, e.g. a rapamycin compound in an emulsion preconcentrate or microemulsion preconcentrate for oral administration. The carrier medium for the rapamycin compound includes a hydrophilic phase, a lipophilic phase and a surfactant. The composition is stable and provides high absorption efficiency. The hydrophilic component, which, with co-components, if present, comprises 10 to 50% by weight of the carrier medium and which is Transcutol, Glycofurol, 1,2-propylene glycol, or mixtures thereof. The lipophilic component, which comprises 10 to 85% by weight of the carrier medium and which is selected from the group consisting of 1) fatty acid triglycerides, 2) mixed mono-, di-, and tri-glycerides, and 3) transesterified ethoxylated vegetable oils. The surfactant, which comprises 5 to 80% by weight of the carrier medium and which is selected from the group consisting of polyethyleneglycol, natural or hydrogenated castor oils, polyethylene-sorbitan fatty acid esters, polyoxyethylene fatty acid esters, polyoxyethylene-polyoxypropylene co-polymers, and block co-polymers. The relative proportion of the active ingredient and components is selected such that on dilution with water, a microemulsion having an average particle size of less than 150 nanometer is spontaneously formed.

Mulye teaches in US patent 6,057,289 about a self-emulsifying pharmaceutical compositions containing an effective amount of cyclosporin in association with a pharmaceutical carrier comprising (a) a cyclosporin solubilizing agent consisting essentially of a solubilizing effective amount of a fatty acid of 6 to 22 carbon atoms, and (b) a non-ionic surfactant having an HLB value greater than 10, said non-ionic surfactant being present with

the cyclosporin solubilizing agent and cyclosporin in an amount sufficient to form an emulsion when brought into contact with an aqueous medium in a mammal. Certain specific compositions may form microemulsions upon mixing with aqueous medium.

Similarly, Lacy et al. teach in US patent 6,096,338 (and in an earlier US patent 5,645,856) to prepare a carrier for hydrophobic drugs, and pharmaceutical compositions based thereon, which carrier comprises a digestible oil and a pharmaceutically acceptable surfactant component for dispersing the oil in vivo upon administration of the carrier, which comprises a hydrophilic surfactant, said surfactant component being such as not to substantially inhibit the in vivo lipolysis of the digestible oil.

The following should be noted about compositions of US patents 6,096,338 and 5,645,856:

- (1) that this carrier system is suitable only for the hydrophobic drugs;
- (2) that the carrier system is designed to facilitate only a digestible oil vehicle;
- (3) on addition into aqueous medium this mixture disperses into an oil-in-water type emulsion;
- (4) that this carrier system does not attempt to obtain a self-dispersible system that results in a micron or sub-micron size surface-modified and surface-stabilized solid particles of the active substance;
- (5) that the surfactant system is designed to avoid inhibitory interference on the gastrointestinal lipolysis of the oil, which is considered by the author as necessary for bioavailability improvement of the orally administered drug; and
- (6) that the compositions of this patent are limited to only the peroral dosage forms.

While it is not intended to present a comprehensive review of background materials, the above examples demonstrate nonexistence of compositions capable of delivering water-insoluble or poorly water-soluble biologically active substances (that may be either oil-soluble or oil-insoluble) via self-dispersible systems providing stable surface-modified micron or sub-micron size solid particles of the active agent. The examples mentioned as the background material, although related to the subject matter of the current inventive compositions and processes, they are different from the present invention.

SUMMARY OF THE INVENTION

This invention disclosure relates to compositions useful for the delivery of stable surface modified sub-micron and micron sized particles of water-insoluble biologically active substances from a carrier system comprising a non-aqueous medium(s) and surfactant(s) and optionally a hydrophilic component that self-disperses on exposure to an aqueous environment. In the aqueous environment the dispersed phase comprises components of the carrier system and the surface modified particles. Surface modified particles refers to micron or sub-micron size solid particles of a water-insoluble active ingredient that is surface-modified by the application of surface active agents to stabilize the said particles against particle size growth, flocculation, aggregation, or agglomeration. In the current invention, the stable surface-modified particulate dispersion of the water-insoluble biologically active substances may be formed in the carrier system using a high-energy dispersion technique such as homogenization or milling. Alternatively, the water-insoluble biologically active substances may be dissolved in a suitable carrier system and then on exposing this solution *in-vitro* or *in-vivo* to aqueous media a self-dispersion of surface modified particles of the water-insoluble drug and other components of the carrier medium is formed. The dispersed surface modified particles may be present in the dispersed non-aqueous medium or in the aqueous phase or both.

The term "water-insoluble biologically active substance" as used herein shall mean any water-insoluble or poorly water-soluble biologically active substance.

The term "non-aqueous composition" as used herein shall refer to any composition that is devoid of water.

The term "carrier system" as used herein is to be understood as defining the material in which the preparation of the stable water-insoluble biologically active substance of this invention is made. The carrier system comprises at least one component from (i) non-aqueous media in which a water-insoluble biological agent is soluble, or non-aqueous media in which a water-insoluble biological agent is insoluble or poorly soluble; and at least one component from (ii) a surfactant system; and optionally at least one component from (iii) a hydrophilic component.

The term "aqueous medium" is meant water; buffered water including phosphate buffered water, phosphate buffered saline, citrate buffered water, acetate buffered water, water buffered with pharmaceutically acceptable pH controlling agents; water containing salts such as sodium chloride and other pharmaceutically acceptable salts; water containing soluble

agents for lyoprotection or cryoprotection such as dextrose, mannitol, trehalose, sucrose, sorbitol, and other pharmaceutically acceptable lyoprotectants and cryoprotectants; water containing soluble agents used to facilitate spray drying such as polyhydroxy-containing compounds such as sugars, polyols, and water containing mixtures of these buffers, agents and compounds. In one embodiment, the aqueous medium can contain one or more soluble surface active agent. In another embodiment, the aqueous medium can contain one or more surfactants dispersed such as by shear mixing into water or other aqueous medium as described herein. In yet another embodiment, the aqueous medium can comprise a biological fluid such as blood, plasma, saliva, urine, a protein-containing solution, an aqueous suspension of a protein, lymph fluid, semen, vaginal fluid, lachrymal fluid, nasal fluid, synovial fluid, cerebral fluid, cerebrospinal fluid, amniotic fluid, pancreatic fluid, pulmonary fluid, ascites fluid, fluid from a cyst, gastric fluid, intestinal fluid, and other fluids that can be removed from a patients for use in this invention. Fluids from an individual patient are inherently biocompatible in most cases with that individual, and a suspension of this invention in a fluid from a patient can be administered to that patient as a biocompatible fluid suspension. Optionally, such patient derived biological fluids can be diluted, for example with other aqueous media as described herein. Optionally, such patient derived biological fluids can be concentrated by removal of some of the water contained therein. Optionally, such patient derived biological fluids can be mixed or blended with other biological fluids from the same patient or from different patients.

In preferred compositions of this invention the non-aqueous carrier system consists of at least one non-aqueous medium component, at least one surfactant component, and optionally at least one hydrophilic component.

Preferred non-aqueous media of the carrier system include hydrophobic components such as triglycerides, diglycerides, monoglycerides, free fatty acids, and fatty acid esters and derivatives thereof, individually or in combination. Examples of such hydrophobic components include but are not limited to propylene glycol dicaprylate/caprate, capric/capric triglyceride, caprylic/capric/linoleic triglyceride, e.g. synthetic medium chain triglycerides having C₈₋₁₂ fatty acid chains or other derivatized (synthetic) triglycerides of the type known and commercially available under Miglyol 810, 812, 818, 829 and 840, linoleic acid, linoleic acid ethyl ester, fish oils as free fatty acids, their esterification and their transesterification products, e.g. of the type known and commercially available under EPAX6000FA, EPAX4510TG, individually or in combination. Additional examples include vegetable oils and C₁₂₋₁₈ fatty acid mono-, di- and triglycerides prepared by individual admixing or as transesterification

products of vegetable oils (such as soybean oil, almond oil, sunflower oil, olive oil or corn oil) with glycerol.

Additional non-aqueous media include pharmaceutically acceptable non-aqueous solvents in which biological substance is either soluble or insoluble or poorly soluble. These media may be derived from the classes of organic chemicals, such as, monohydric alcohols *e.g.*, alkanols; dihydric alcohols *e.g.*, glycols; polyhydroxy compounds *e.g.*, glycerin; aromatic esters, *e.g.*, benzyl benzoate, diethyl phthalate, propyl gallate, triacetin, diacetin, monoacetin, triethyl citrate; water soluble organic solvents such as propylene carbonate and glycofurol, dimethyl isosorbide, dimethyl isoidide and dimethyl isomannide.

One or more surfactants utilized in the carrier of this invention is termed as the surfactant system. In those compositions possessing more than one surface active agents, the principle surface active agent, that is present in larger quantity is called the surfactant and the other surface active agents are named as co-surfactants.

The surfactant system of this invention may include natural or synthetic surface modifying agents selected from phospholipids, and non-ionic, anionic or cationic surfactants or combinations thereof. These surface modifying molecules adhere onto the surfaces of the insoluble-particles of biologically active substances and stabilize the sub-micron and micron size drug particles. The stabilization of the particulate matter against aggregation, flocculation, or particle-size growth is thought to arise from lipophilic, amphipathic, and/or electrostatic interactions as well as steric hindrance. In addition, alteration of the density of the medium, the polarity and the surface tension of the particles suspended in the medium, and favorable adjustment of the hydrodynamic properties reduce the tendency of dispersed particles to either cream or sediment. Also, addition of the surface-active substances to these compositions enables them to easily disperse upon release into an aqueous environment such as a biological fluid, particularly gastric and gastro-intestinal fluids, blood, and lymph.

In the compositions of this invention a preferred surfactant system comprises at least one phospholipid selected from the group consisting of saturated phospholipids, unsaturated phospholipids, synthetic phospholipids, natural phospholipids, and combinations thereof.

The hydrophilic component when optionally used comprises less than about 10% of the carrier system. Examples of hydrophilic components include low-molecular weight monohydric alcohols and preferably ethanol, low-molecular weight polyhydric alcohols,

glycols, and glycerol, and mixtures thereof. In a preferred embodiment, the hydrophilic component comprises a pharmaceutically acceptable monohydric or polyhydric alcohol.

A typical carrier system of this invention consists of one or more suitable hydrophobic oils and a surfactant system comprising at least one surface modifying agent and optionally one or more hydrophilic components. The composition of this invention may remain in a solid state or a semi-solid state such as waxy or gel state or in liquid state. In addition to the carrier system the composition of this invention may also preferably additionally comprise further adjuvants like anti-oxidants, preserving agents and stabilizers, flavoring agents, thickening agents, diluents and other pharmaceutically acceptable ingredients for other purposes. The compositions of this invention are highly concentrated and suitable for packaging as a capsule or tablet and thus are very convenient for use in oral dosage forms. Additionally, compositions of this invention may be used to effectively deliver biologically active substances via other routes of drug delivery, such as, but not limited to peroral, parenteral, transdermal, pulmonary (inhalation), ophthalmic, transmucosal routes or by injection.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a schematic illustration of surface modified particles of water-insoluble substance in non-aqueous medium dispersed in aqueous medium.

Figure 2 is a process flowchart for obtaining micronized and surface stabilized biological substance dispersed in non-aqueous self-dispersing carrier medium having particles in a size range of about 0.01 to about 10 micrometer, and preferably 0.01 to 2 micrometer.

Figure 3 is a process flowchart for obtaining anti-solvent precipitated and surface stabilized biological substance dispersed in non-aqueous self-dispersing carrier medium having particles in a size range of about 0.01 to about 10 micrometer, and preferably 0.01 to 2 micrometer.

Figure 4 is a process flowchart for obtaining dispersion of surface stabilized biological substance in non-aqueous self-dispersing carrier medium having particles in a size range of about 0.01 to about 10 micrometer, and preferably 0.01 to 2 micrometer.

Figure 5 is a process flowchart for obtaining a solution of water-insoluble active substance in non-aqueous self-dispersing carrier medium.

DESCRIPTION OF THE INVENTION

Surprisingly, it has been found that suspensions of stable surface modified micron and sub-micron sized particles in self-dispersing carrier systems are attainable. These particles consist of a water-insoluble biologically active substance coated with at least one surface-modifying agent, *e.g.*, phospholipid molecules or other surface modifying agents or combinations thereof. Preferably the particles are coated with phospholipid(s).

In one embodiment of this invention particles of a biologically active water-insoluble or poorly soluble substance of about 0.01 to about 10 micrometer volume weighted mean particle size are stabilized with surface modifiers and dispersed in a non-aqueous medium in which the substance is insoluble or poorly soluble. The active substance may be released to an aqueous environment as surface modified submicron or micron sized particles of the water-insoluble substance from this composition.

In another embodiment of this invention the above mentioned composition of the water-insoluble biologically active substance is added to an aqueous medium, for example, the fluid medium of an *in-vivo* environment, and self-disperses to form non-aqueous droplets and surface modified particles (see Figure 1). The active substance may be released to the *in-vivo* environment from this self-dispersion.

Figure 1 is a schematic illustration of surface modified particles of water-insoluble substance in non-aqueous medium dispersed in aqueous medium. Item 10 represents the aqueous environment or aqueous medium. Item 20 represents the self-dispersed non-aqueous medium droplet. Item 30 represents stable surface modified particles of water-insoluble or poorly water-soluble biologically active substance. Item 31 represents surface modified particles of water-insoluble or poorly water-soluble biologically active substance migrated to the aqueous medium. Item 32 is a magnified schematic illustration of the surface modified particles of water-insoluble or poorly water-soluble biologically active substance.

In another embodiment of this invention, there is provided composition comprising stable particles of a surface modified water-insoluble biologically active substance of a mean size in the range of 0.01 to 10 micrometers, which particles are dispersed in a non-aqueous carrier system comprised of a non-aqueous medium in which the biologically active substance is not soluble or is poorly soluble; and a surfactant system consisting of at least one surfactant which is soluble in the non-aqueous medium and which adsorbs to the surface of the biologically active substance; and optionally a quantity of not more than about 10% of the total

weight of said composition of one or more hydrophilic substance that provides a self-dispersing property to said composition wherein said composition can self-disperse upon addition to an aqueous medium to form a suspension of components of the non aqueous carrier system and stable particles of said water-insoluble biologically active substance and wherein said particles have a size in the range of 0.01 to 10 micrometer and have associated therewith on the surface at least a portion of said surfactant system.

In yet another embodiment of the invention there is provided composition comprising a water-insoluble biologically active substance mixed with a non-aqueous carrier system, wherein said carrier system comprises a non-aqueous medium in which said water-insoluble biologically active substance is soluble; and a surfactant system consisting of at least one surfactant; and optionally a quantity of not more than about 10% of the total weight of said composition of one or more hydrophilic substance that provides a self-dispersing property to said composition; wherein said composition can self-disperse upon addition to an aqueous medium to form a suspension comprising components of the non-aqueous carrier system and stable particles of said water-insoluble biologically active substance wherein said particles have a mean size in the range of 0.01 to 10 micrometer and have associated therewith on the surface at least a portion of said surfactant system.

Surface modifiers are selected such that they adsorb to the surface of the water-insoluble substance when exposed to the non-aqueous carrier and when dispersed in aqueous media to allow the formation of stable small particles of the substance (Figure 1).

In yet another embodiment of the invention, the carrier system consists of non-aqueous medium and a surfactant system in which a biologically active substance is dissolved, such that upon mixing with an aqueous environment the composition self-disperses to form an oil-in-water dispersion in which the droplet size is sufficient to entrap one or more particles of biologically active substance. Alternatively, upon mixing with an aqueous environment the composition may self-disperse to form surface modified micron and submicron sized particles of the water-insoluble substance wherein the particles are at least in-part external to the non-aqueous carrier system.

Such novel compositions provide preferred dosage forms of pharmaceutical agents useful to achieve certain pharmacological actions such as controlled and/or sustained delivery of a drug substance, targeted delivery of a drug substance such as to the lymphatic system. They also can provide a dosage form with reduced tissue irritation such as commonly found after oral delivery of non-steroidal anti-inflammatory drugs.

In addition, there are applications where aqueous medium may not be suitable for the surface modified sub-micron particulate suspensions. Presence of water in these suspensions may not allow the use of certain packaging systems such as hard- or soft-gelatin capsules or starch capsules because these packaging systems are known to dissolve in aqueous media. Preferred medium for such non-aqueous particulate suspensions or dispersions may be a suitable oil or a suitable hydrophobic liquid or a suitable amphipathic medium to allow a traditional packaging system, such as a tablet, gelcap, or capsule to be employed.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention can be practiced with a wide variety of water-insoluble biologically active substances that are either insoluble or poorly soluble in hydrophobic liquids. One embodiment of this invention may utilize even those water-insoluble biologically active substances that are soluble in non-aqueous media.

Examples of some preferred biologically active agents useful in this invention include antifungal agents, immunosuppressive and immunoactive agents, antiviral agents, antineoplastic agents, analgesic and antiinflammatory agents, antibiotics, antiepileptics, anesthetics, hypnotics, sedatives, antipsychotic agents, neuroleptic agents, antidepressants, anxiolytics, anticonvulsant agents, antagonists, neuron blocking agents, anticholinergic and cholinomimetic agents, antimuscarinic and muscarinic agents, antiadrenergic and, antarrhythmics, antihypertensive agents, antineoplastic agents, hormones, nutrients, oligonucleotides, vaccine, peptides or peptidomimetics and proteins. A detailed description of these drugs may be found in *Remington's Pharmaceutical Sciences*, 18th Edition, 1990, Mack Publishing Co., PA.

More preferred biologically active substances include nifedipine, alfaxalone, budesonide, a hormone, ursadiol, propranolol, itraconazole, acyclovir, fenofibrate and their analog derivatives. Potential imaging agents include water-insoluble X-ray or Magnetic Resonance Imaging nanoparticulate material.

In one aspect, preferred biologically active substances members of the group consisting of an antihypertensive drug; nifedipine; an anticholinergic drug; ursodiol; a drug for treating a gastro-intestinal disorder; budesonide; an antineoplastic drug; peclitaxel; camptothecin; a derivative of peclitaxel; a derivative of camptothecin; an NSAID; piroxicam; an anti-fungal agent; itraconazole; an anti-viral agent; acyclovir; a derivative of acyclovir; a cholesterol

controlling agent; fenofibrate; an immuno-suppressive peptide; cyclosporine; a protein used in the treatment of diabetes; insulin; and a derivative of insulin.

Preferred non-aqueous media of the carrier system include hydrophobic components such as triglycerides, diglycerides, monoglycerides, saturated or unsaturated free fatty acids, mixtures of saturated and unsaturated free fatty acids, and fatty acid esters and derivatives thereof, individually or in combination. Examples of such hydrophobic components include but are not limited to propylene glycol dicaprylate/caprate, caprilic/capric triglyceride, caprylic/capric/linoleic triglyceride, e.g. synthetic medium chain triglycerides having C_{8-12} fatty acid chains or other derivatized (synthetic) triglycerides of the type known and commercially available under Miglyol 810, 812, 818, 829 and 840, as well as linoleic acid, linoleic acid ethyl ester, fish oils as free fatty acids, their esterification and their transesterification products, e.g. of the type known and commercially available under EPAX6000FA, EPAX4510TG, all individually or in combination. Additional examples include vegetable oils and C_{12-18} fatty acid mono-, di- and triglycerides prepared by individual admixing or as transesterification products of vegetable oils (such as soybean oil, almond oil, sunflower oil, olive oil or corn oil) with glycerol.

Additional non-aqueous media include pharmaceutically acceptable non-aqueous media or solvents in which the biological substance is either soluble or insoluble or poorly soluble. These solvents may be selected from classes of organic chemicals such as but not limited to, monohydric alcohols e.g., alkanols; dihydric alcohols e.g., glycols; polyhydroxy compounds e.g., glycerin; aromatic esters, e.g., benzyl benzoate, diethyl phthalate, propyl gallate; non-aromatic esters such as triacetin, diacetin, monoacetin, triethyl citrate; water soluble organic solvents such as propylene carbonate and glycofurol, dimethyl isosorbide, dimethyl isoside and dimethyl isomannide.

Surface modifiers or their combinations are selected from known organic and inorganic excipients suitable for the delivery of the biologically active substance such as pharmaceutical excipients. Such excipients include various pharmaceutically useful polymers, low molecular weight oligomers, natural products and surfactants. Preferred surface modifiers include nonionic and ionic surfactants. Examples of some suitable surface modifiers are:

- (a) natural or synthetic lipophilic agents, e.g., phospholipids, cholesterol, and cholesteryl fatty acid esters and their surface active derivatives;

- (b) nonionic surfactants such as polyoxyethylene fatty alcohol ethers, sorbitan fatty acid esters, polyoxyethylene sorbitan fatty acid esters (Tweens), sorbitan esters, glycerol esters such as Myrj and glycerol triacetate (triacetin), polyethylene glycols, cetyl alcohol, cetostearyl alcohol, stearyl alcohol, poloxamers, polaxamines, polyoxethylene castor oil derivatives (Cremophors), vitamin E or its derivatives, such as D-a-tocopheryl polyethylene glycol 1000 succinate (vitamin E TPGS), PEG glyceryl fatty acid esters such as PEG-8 glyceryl caprylate/caprate (commercially known as Labrasol), PEG-4 glyceryl caprylate/caprate (Labrafac Hydro WL 1219), PEG-32 glyceryl laurate (Gelucire 44/14), PEG-6 glyceryl mono oleate (Labrafil M 1944 CS), PEG-6 glyceryl linoleate (Labrafil M 2125 CS), propylene glycol mono- and di-fatty acid esters, such as propylene glycol laurate, propylene glycol caprylate/ caprate; also diethylene glycol monoethyl ether, commercially known as transcutol, sorbitan fatty acid esters, such as the type known and commercially available under the trade name Span (e.g., Span 20); monoglycerides and acetylated monoglycerides, e.g., glycerol monooleate, glycerol monostearate and mono-and di-acetylated monoglycerides, monoacetin, diacetin; carbomers, such as the type known and commercially available under the trade name Carbopol;
- (c) anionic surfactants such as bile salts, potassium laurate, triethanolamine stearate, sodium lauryl sulfate, alkyl polyoxyethylene sulfates, sodium alginate, dioctyl sodium sulfosuccinate, sodium carboxymethylcellulose, and calcium carboxymethylcellulose;
- (d) cationic surfactants such as quaternary ammonium compounds, benzalkonium chloride, cetyltrimethylammonium bromide, and lauryldimethylbenzyl-ammonium chloride;
- (e) substituted cellulose derivatives such as methylcellulose, hydroxycellulose, hydroxy propylcellulose, hydroxy propylmethylcellulose, noncrystalline cellulose, sodium carboxymethyl cellulose;
- (f) polyethylene glycol (PEG); e.g. PEG 300, PEG 400, PEG 600, PEG 1000, PEG 1500, PEG 3400; such as the type known and commercially available under the trade name Carbowax, Lutrol E and Hodag PEG;
- (g) colloidal clays such as bentonite and veegum;
- (h) natural proteins such as casein, gelatin;
- (i) trigacanth, waxes, enteric resins, paraffin, acacia.

The non-aqueous hydrophobic media are selected from triglycerides, oils derived from vegetable or animal origins such as vegetable oils or fish oils, free fatty acids such as oleic or linoleic acids or other poly-unsaturated fatty acids, fatty acid esters, and other pharmaceutically suitable hydrophobic compounds that remain in liquid state at ambient temperature and pressure such as hydrofluorocarbons such as perflubron.

Optionally, small quantity of low-molecular weight hydrophilic substances, for example, monohydric or polyhydric alcohols, such as ethanol or glycols or glycerol may be also added.

A detailed description of these surface modifying agents and lipophilic or hydrophobic media may be found in *Remington's Pharmaceutical Sciences*, 18th Edition, 1990, Mack Publishing Co., PA; and *Theory and Practice of Industrial Pharmacy*, Lachman *et al.*, 1986.

In compositions of this invention possessing more than one surface active agent the principle surface active agent, *i.e.*, that present in largest quantity is called the surfactant and the other surface active agents are called as co-surfactants. When present in equal amounts, the surfactant is defined by the higher molar amount if w/w ratio is used or the higher weight amount if molar ratio is used.

The type and amount of surfactant and co-surfactant used is based on the relative solubility and/or polarity of these ingredients. The formulation compositions are hence optimized with respect to each biologically active agent individually. It is thought that some of the functions of the surface modifier(s) as it relates to this invention are:

- to suppress the process of Ostwald Ripening and therefore maintain the particle size, increase the storage stability, minimize sedimentation, and decrease the particle growth during lyophilization and other unit-processes of the manufacture of these compositions;
- adhere or coat firmly onto the surfaces of particles and therefore modify the interfaces between the particles and the non-aqueous carrier system and the aqueous environment when self-dispersion occurs;
- increase the interface compatibility between particles and the non-aqueous carrier system and the aqueous environment when self-dispersion occurs; and
- facilitate self-emulsification or self-dispersion capability of the composition.

METHODS

A dispersion of particles of a biologically active substance in a non-aqueous carrier system in which it is insoluble or poorly soluble is formed using one of the following four processes. These processes are schematically outlined in Figures 2-5.

Process I: Micronized surface stabilized water-insoluble biological substance dispersed in a non-aqueous carrier system.

The first of these processes (Figure 2) utilizes a particle size reduction unit process to form aqueous suspensions of surface modified sub-micron or micron sized particles of the water-insoluble substances that are also insoluble in the non-aqueous carrier system. In this process an aqueous medium premix of the water-insoluble biologically active substance and the surfactant system is prepared and then subjected to one of the particle size reduction unit processes. A suspension of surface modified small particles, ranging from 0.01 to 10 micrometer volume weighted mean particle size is obtained. This suspension is subjected to a drying unit process such as spray drying or lyophilization yielding a dry powder of the particles, which is then mixed or homogenized with or without high shear with non-aqueous carrier system, such as oily medium, and optionally with additional components of the surfactant system. The dispersion of the particles in the self-dispersing non-aqueous carrier so obtained is then available for final packaging.

Process II: Anti-solvent precipitated surface stabilized water-insoluble biological substance dispersed in non-aqueous carrier.

In the second process (Figure 3) the water-insoluble biologically active substance is dissolved in appropriate organic solvent. Surface modifier of the surfactant system is mixed either in this solution or in an antisolvent for the biologically active substance, which can be an aqueous medium. The antisolvent medium and the organic solution with the biologically active substance are mixed to cause the antisolvent precipitation of the biologically active substance. If required, this dispersion is further subjected to a size reduction unit process (described below). A suspension of surface modified small particles, ranging from 0.01 to 10 micrometer volume weighted mean particle size is obtained. This suspension is subjected to a solvent and antisolvent removal unit process such as spray drying or lyophilization yielding a dry powder of the surface modified particulate composition, which is then mixed or homogenized with or without high shear with the non-aqueous carrier system, such as the oily medium, and

optionally with additional components of the surfactant system. The non-aqueous dispersion of the microparticulate suspension so obtained is then available for final packaging.

Process III: Dispersion of surface stabilized water-insoluble biological substance in non-aqueous self-emulsifying carrier system.

In the third process (Figure 4) is performed a particle size reduction unit process (see below) on the premix of water-insoluble biologically active substance and the carrier system to directly produce the suspensions of the active substances that are insoluble or poorly-soluble in the carrier system. The non-aqueous dispersion of the particles in the self-dispersing non-aqueous carrier system so obtained is then available for final packaging.

In this case, the composition is selected such that upon addition of aqueous medium these compositions self-disperse. The droplets of the self-dispersion contain particles of the surface stabilized water-insoluble biological substance suspended in the oily droplets of the dispersion.

Process IV: Dispersion of solution surface stabilized water-insoluble biological substance in non-aqueous self-emulsifying medium.

The fourth process (Figure 5) relies on making compositions of the substances that are soluble in the carrier system such that upon addition of aqueous medium these compositions self-disperse into sub-micron and micron sized particles of surface stabilized water-insoluble biological substance. The non-aqueous mixture of the biologically active substance so obtained is then available for final packaging.

In this case, the composition is selected such that upon addition of aqueous medium these compositions self-disperse. The droplets of the self-dispersion may contain the surface modified particles of the water-insoluble biologically active substance suspended in the oily droplets of the dispersion. Alternatively, the self-dispersion process may provide a particle suspension of the active substance wherein the components of the carrier system adhere to the small active substance particle to facilitate spontaneous formation of the said dispersed surface modified particles.

Particle Size Reduction Unit Processes:

The particles must be reduced in size at a temperature that does not significantly degrade the drug substance. Processing temperatures of less than about 30-40 degree C are usually preferred. If required, the processing equipment and the process-fluid flow is cooled

with conventional cooling equipment. The method is conveniently carried out under conditions of ambient temperature. When size reduction is carried out in aqueous medium, pH is a very important parameter to control. It is controlled within a range so that the active and the inactive ingredients do not undergo pH-induced degradation. Similarly for the compounds sensitive to oxidation proper nitrogen blanketing procedure is applied.

High Energy Micronization: The biologically active substance together with other appropriate ingredients are subjected to high energy provided by high pressure homogenization or microfluidization as known in the art. In these processes, the high energy results in shear, impaction, cavitation and other forces that causes particle size reduction.

Media Milling

The particle size reduction of the drug substance can be achieved by a dispersion mill. Suitable dispersion mills include a ball mill, an attritor mill, a vibratory mill, and media mills such as a sand mill and a bead mill. The grinding media for the particle size reduction step can be selected from rigid media preferably spherical or particulate in form having an average size less than about 3 mm and, more preferably, less than about 1 mm. The selection of material for the grinding media is not believed to be critical. We have found that zirconium oxide, such as zirconia stabilized with magnesia, zirconium silicate, stainless steel, titania, alumina, zirconia stabilized with yttrium and glass grinding media provide particles having minimal levels of contamination which are believed to be acceptable for the preparation of pharmaceutical compositions.

Rapid Expansion of a Supercritical Solution

The biologically active substance is dissolved in a supercritical gas such as carbon dioxide that may also contain the surface modifiers and then the solution is expanded over an aqueous suspension with or without the surface modifiers. The principle feature of this unit process is rapid attainment of intimate contact of the dissolved drug and the surface modifier during the very fast precipitation step of the drug from their solution in the liquefied gas. While very rapid precipitation is a characteristic of precipitation of solutes from supercritical fluids, the rapid intimate contact with the surface modifier is achieved by having the surface modifiers dissolved in the liquefied-gas containing the dissolved drug. A rapid intimate contact between the surface modifier and the newly formed particle substantially inhibits the crystal growth of the newly formed particle.

Although at least one (first) surface modifier should be dissolved along with the water insoluble substance to be reduced in size in the liquefied gas, additional (second) surface modifying agents of the same or different chemical nature may also be included in the aqueous medium. Further, during or after precipitation the fluid streams may be subjected to additional high shear forces, cavitation or turbulence by a high-pressure homogenizer to facilitate intimate contact of the particle surface and the surface modifier. Thus, in those cases where all the surface modifier is dispersed in the aqueous medium and the liquefied gas contains only the water insoluble substance, additional high shear forces, cavitation or turbulence by a high-pressure homogenizer can be exploited to facilitate the intimate contact of the particle surface and the surface modifier.

Antisolvent Precipitation

A solution of the biologically active substance and other ingredients in a volatile solvent is mixed with an antisolvent which cannot solubilize the formulation components, and thus results in the active substance precipitated together with other ingredients. Suitable particle size distribution of precipitate is achieved by appropriate control of the conditions such as concentration, rate of mixing of the solution and anti-solvent, temperature and pressure. Solvent or antisolvent may be selected from suitable organic solvents as well as supercritical fluids, for instance, liquefied gasses or supercritical fluids such as supercritical carbon dioxide.

Liquid or Solvent or Antisolvent Media Removal:

Excess liquid dispersing medium that contains the micronized particles, such as water or organic solvents, is removed by the process of drying. In case the solvent or anti-solvent medium is a supercritical compressed fluid, it is removed by decompression. The decompression of supercritical compressed fluid solution also results in the microparticulate formation.

EXAMPLES

Example 1: Solubility of biologically active substances, particularly drug substances in lipophilic and hydrophilic solvents. The example drug, Itraconazole was found to be very poorly soluble in pharmaceutically acceptable lipophilic vehicles. Itraconazole solubility of at least 25 mg/mL in oil is required for making an oil-in-water type emulsion-formulation. The solvents mentioned in the following table were evaluated. Solubilities, determined by high performance liquid chromatography (HPLC) assay of the dissolved drug in water and several

non-aqueous vehicles, are given the following table in units of mg/mL. Low solubility of this drug in either water or organic solvents renders it suitable for formulating as a stable surface modified particulate dispersion in a non-aqueous carrier system.

Solubility of Itraconazole (mg/mL) in some Solvents by HPLC

Water	0.01
Decyl oleate	0.02
Ethyl oleate	0.04
Ethyl myristate	0.06
Isopropyl myristate	0.08
Ethyl caprate	0.10
Miglyol-840	0.20
Soybean oil	0.25
Miglyol 810	0.34
Capric triglyceride	0.37
Ethyl alcohol	0.54
Corn oil PEG-6 ester	1.24
Propyleneglycol laurate	1.26
Ethyl caprylate	1.57
Miglyol 818	1.59
Apricot kernel oil	2.65
Linoleic acid	2.73
PEG-200	3.05
PEG-300	3.26
PEG-400	3.33
Triethyl citrate	4.12
Miglyol 812	5.14

Glycerol triacetate	7.55
Glycerol a,a'-diacetate	9.11
1,2 Propane diol	12.5
Glyceryl linoleate	13.7
Plurol oleique CC 497	16.0

Example II: Solubilization and precipitation behavior of some drugs. As an example, 20 mg of either nifedipine or ursadiol were added into 1 mL of either a medium chain triglyceride, Miglyol 810, or to triacetin, or to benzyl benzoate. Almost all of the nifedipine that was added was observed to remain unsolubilized. Similarly ursodiol was not soluble in either Miglyol 810 or triacetin. This makes it possible to prepare compositions of nifedipine and ursodiol in these non-aqueous media as microparticulate suspension by further particle size reduction.

However, ursodiol was found to dissolve in benzyl benzoate. A portion of the ursodiol solution in benzyl benzoate was transferred to water and shaken. A mixture containing oily droplets resulted. No precipitation was seen occurring. On examining the mixture by an optical microscope under polarized light conditions surprisingly particles of regular crystalline shape and below 10 micron size were observed. This observation makes it possible to further prepare compositions of ursodiol utilizing appropriate solvent system and surface active agents so that micron-sized particle will be generated spontaneously upon addition of these compositions in aqueous media.

Example III: A mixture containing itraconazole and egg-phospholipid in ethyl oleate was subjected to high-pressure homogenization in an Avestin Emulsiflex C5 homogenizer. Homogenizing pressures of up to 20,000 psi were used. The process fluid temperature was maintained at below 40°C by cooling with a heat exchanger equipped with a jacket of coolant fluid. Fine particulate suspension of volume weighted mean particle size of about 0.6 micron was produced.

Example IV: Many compositions containing cyclosporine, omega-3 fatty acid, its ethyl ester or triglyceride, surfactants such as vitamin E-TPGS, Tween-20, Tween-80, Labrasol, Pluronic F68, Pluronic L44, co-surfactants such as ethanol, propylene glycol, or sodium

hydroxide dissolved in either ethanol or propylene glycol were prepared. These compositions were highly concentrated and could be encapsulated in appropriate capsules. Upon adding to aqueous medium many of these compositions produced opaque milky-white suspensions containing surface-modified micron or sub-micron size solid particulate suspension. Thus these compositions can be further developed in the ingestible dosage-forms that facilitate efficient delivery of the drug substance upon mixing with aqueous media within *in-vivo* environment via surface-modified micron or sub-micron size particulate suspension.

Examples of cyclosporine and fenofibrate compositions which self-disperse into micron or sub-micron size solid particulate suspensions on addition of aqueous medium are shown in Tables I, II and III. Required amounts (in milligrams) of the ingredients were weighed into a 20-mL clear glass vial and mixed until complete dissolution was observed. Twenty milliliters water or simulated gastric fluid (SGF) or SGF mixed with water was added and gently mixed by inverting the mixture by hand several times. Milky white microparticulate suspension formed and demonstrated self-dispersibility property of the composition.

Table I: Examples of the compositions of cyclosporine that self-disperse into surface-modified micron or sub-micron sized particle suspensions. Milligrams of the components used are shown.

FORMULATION ID →	1.1	1.2	1.3	1.4
COMPOSITION ↓				
Oil Phase				
Cyclosporine	50	50	50	50
EPAX4510TG	150	150	150	150
Surfactants				
Vit-E-TPGS	45	37	30	22
Tween-80	405	338	270	203
Co-Surfactants				
Ethanol	150	150	150	150

Table III: Examples of fenofibrate compositions that self-disperse into surface-modified micron or sub-micron sized particulate suspensions. Milligrams of the components used are shown.

Formulation ID	Miglyol 810	Linoleic acid	Fenofibrate	Tween 80	Vitamin E-TPGS	Ethanol	Propylene Glycol
3.1	260.0	130.0	100.0	450.0		125.0	124.0
3.2	260.0	130.0	100.0	375.0	76.0	125.0	124.0
3.3	260.0	130.0	100.0	436.0	88.0	133.0	48.0
3.4	260.0	130.0	100.0	400.0	74.0	188.0	65.0
3.5	260.0	130.0	100.0	436.0	88.0	89.0	89.0
3.6	260.0	130.0	100.0	400.0	75.0	125.0	

Compositions in accordance with the present invention may include additional ingredients, for example, diluents or bulking agents, anti-oxidants, preserving agents, and pH buffering agents. The above examples are illustrative but not limiting examples of compositions in accordance with the present invention. The invention has been described with reference to the preferred embodiments thereof, but variations and modifications can be affected within the spirit and scope of the invention.

WHAT IS CLAIMED IS:

1. A composition comprising stable particles of a surface modified water-insoluble biologically active substance of a mean size in the range of 0.01 to 10 micrometers, which particles are dispersed in a non-aqueous carrier system comprised of:
 - (a) a non-aqueous medium in which the biologically active substance is not soluble or is poorly soluble; and
 - (b) a surfactant system consisting of at least one surfactant which is soluble in the non-aqueous medium and which adsorbs to the surface of the biologically active substance; and
 - (c) optionally a quantity of not more than about 10% of the total weight of said composition of one or more hydrophilic substance that provides a self-dispersing property to said composition.

wherein said composition can self-disperse upon addition to an aqueous medium to form a suspension comprising components of the non aqueous carrier system and stable particles of said water-insoluble biologically active substance wherein said particles have a size in the range of 0.01 to 10 micrometer and have associated therewith on the surface at least a portion of said surfactant system.

2. A composition comprising a water-insoluble biologically active substance mixed with a non-aqueous carrier system, wherein said carrier system comprises:
 - (a) a non-aqueous medium in which said water-insoluble biologically active substance is soluble; and
 - (b) a surfactant system consisting of at least one surfactant; and
 - (c) optionally a quantity of not more than about 10% of the total weight of said composition of one or more hydrophilic substance that provides a self-dispersing property to said composition;

wherein said composition can self-disperse upon addition to an aqueous medium to form a suspension comprising components of the non aqueous carrier system and stable particles of said water-insoluble biologically active substance wherein said particles

have a mean size in the range of 0.01 to 10 micrometer and have associated therewith on the surface at least a portion of said surfactant system.

3. The compositions of claim 1 and 2 where the non-aqueous carrier system consists of:
 - (a) at least one non-aqueous medium component selected from the group consisting of oils derived from vegetable or animal origins; vegetable oils; fish oils; fish oil free fatty acids; oleic acid; linoleic acid; poly-unsaturated fatty acids; fatty acid esters; triglycerides; caprylic/capric triglyceride; caprylic/capric/linoleic triglyceride; synthetic medium chain triglycerides having C₈₋₁₂ fatty acid chains; synthetic triglycerides; Miglyol 810, Miglyol 812, Miglyol 818, Miglyol 829, Miglyol 840; diglycerides; monoglycerides; monoglyceride and diglyceride free fatty acids; fatty acid esters; propylene glycol dicaprylate/caprate; linoleic acid ethyl ester; EPAX6000FA; EPAX4510TG; cholesteryl fatty acid esters; C₁₂₋₁₈ fatty acid monoglycerides, C₁₂₋₁₈ fatty acid diglycerides, and C₁₂₋₁₈ fatty acid triglycerides prepared from soybean oil, almond oil, sunflower oil, olive oil, and corn oil with glycerol; pharmaceutically acceptable monohydric alcohols; pharmaceutically acceptable alkanols; pharmaceutically acceptable dihydric alcohols; glycols; pharmaceutically acceptable polyhydroxy compounds; glycerin; pharmaceutically acceptable aromatic esters; benzyl benzoate; diethyl phthalate; propyl gallate; triacetin; diacetin; monoacetin; triethyl citrate; water soluble organic solvents; propylene carbonate; glycofurol; dimethyl isosorbide; dimethyl isodide; dimethyl isomannide; pharmaceutically suitable hydrophobic organic solvents; hydrofluorocarbons; and perflubron; and
 - (b) at least one surfactant component selected from the group consisting of natural or synthetic amphiphilic agents; phospholipids; cholesterol; nonionic surfactants; polyoxyethylene fatty alcohol ethers; sorbitan fatty acid esters; polyoxyethylene sorbitan fatty acid esters; Tweens; sorbitan esters; Myrj glycerol esters; glycerol triacetate; triacetin; polyethylene glycols; cetyl alcohol; cetostearyl alcohol; stearyl alcohol; poloxamers; polaxamines; polyoxyethylene castor oil derivatives; Cremophors; vitamin E; D-alpha-tocopheryl polyethylene glycol 1000 succinate; vitamin E TPGS; PEG glyceryl fatty acid esters; PEG-8 glyceryl caprylate/caprate; Labrasol; PEG-4 glyceryl caprylate/caprate; Labrafac Hydro WL 1219; PEG-32 glyceryl laurate; Gelucire 44/14; PEG-6

glyceryl mono oleate; Labrafil M 1944 CS; PEG-6 glyceryl linoleate; Labrafil M 2125 CS; propylene glycol mono fatty acid esters; propylene glycol di-fatty acid esters; propylene glycol laurate; propylene glycol caprylate/caprate; diethylene glycol monoethyl ether; transcutoil; sorbitan fatty acid esters; Span fatty acid esters; Span 20; monoglycerides; acetylated monoglycerides; glycerol monooleate; glycerol monostearate; mono-acetylated monoglycerides; di-acetylated monoglycerides; monoacetin; diacetin; carbomers; Carbopol; anionic surfactants; fatty acid salts; bile salts; potassium laurate; triethanolamine stearate; sodium lauryl sulfate; alkyl polyoxyethylene sulfates; sodium alginate; dioctyl sodium sulfosuccinate; sodium carboxymethylcellulose; calcium carboxymethylcellulose; cationic surfactants; pharmaceutically acceptable quaternary ammonium compounds; benzalkonium chloride; cetyltrimethylammonium bromide; and lauryldimethylbenzylammonium chloride; substituted cellulose derivatives; methylcellulose; hydroxycellulose; hydroxy propylcellulose; hydroxy propylmethylcellulose; noncrystalline cellulose; sodium carboxymethyl cellulose; polyethylene glycol; PEG; PEG 300; PEG 400; PEG 600; PEG 1000; PEG 1500; PEG 3400; Carbowax; Lutrol E; Hodag PEG; and

- (c) optionally at least one hydrophilic component selected from the group consisting of low-molecular weight monohydric alcohols; low-molecular weight polyhydric alcohols; ethanol; glycols; glycerol; and mixtures thereof.
4. Compositions of claim 1 and 2 in which the surfactant system comprises at least one phospholipid selected from the group consisting of saturated phospholipids, unsaturated phospholipids, synthetic phospholipids, natural phospholipids, and combinations thereof.
 5. Compositions of claim 1 and 2 in which the non-aqueous medium comprises an oil selected from the group consisting of a vegetable oil, an animal oil, a synthetic oil, a free fatty acid oil, a poly-unsaturated fatty acid oil, a fatty acid ester oil, a triglyceride oil, a diglyceride oil, a monoglyceride oil, and combinations thereof.

6. The composition of claim 1 where the stable particles are produced in a size reduction unit operation selected from the group consisting of homogenization, milling, microfluidization, precipitation, recrystallization, anti-solvent precipitation, and precipitation from an expanding supercritical fluid.
7. The homogenization unit operation of claim 6 which employs a high pressure homogenizer or microfluidizer selected from the group consisting of an Avestin homogenizer or microfluidizer, a Niro homogenizer or microfluidizer, a Rannie homogenizer or microfluidizer, a Gaulin homogenizer or microfluidizer, a homogenizer or microfluidizer comprising of a homogenizing valve made of ceramic materials, and a homogenizer or microfluidizer comprising of a homogenizing valve made of diamond materials.
8. Compositions of claims 1 and 2 wherein the concentration of the biologically active substance is sufficient for use in sustained or controlled delivery of the active substance.
9. Compositions of claims 1 and 2 further comprising one or more pharmaceutical excipient useful for peroral, parenteral, transdermal, inhalation, or ophthalmic administration of the biologically active substance.
10. Compositions and processes according to claims 1 and 2 wherein the biologically active substance is selected from the group consisting of an antihypertensive drug; nifedipine; an anticholinergic drug; ursodiol; a drug for treating a gastro-intestinal disorder; budesonide; a hormone; an antineoplastic drug; peclitaxel; camptothecin; a derivative of peclitaxel; a derivative of camptothecin; an NSAID; piroxicam; an anti-fungal agent; itraconazole; an anti-viral agent; acyclovir; a derivative of acyclovir; a cholesterol controlling agent; fenofibrate; an immuno-suppressive peptide; cyclosporine; a protein used in the treatment of diabetes; insulin; and a derivative of insulin.

11. The compositions of claims 1 and 2 contained in a capsule selected from the group consisting of a hard gelatin capsule, a soft gelatin capsule, and a starch capsule, which capsule optionally comprises a pharmaceutically acceptable coating for controlling the release of the biologically active substance.
12. The compositions of claims 1 and 2 contained in a tablet, which tablet optionally comprises a pharmaceutically acceptable coating for controlling the release of the biologically active substance.
13. The compositions of claims 1 and 2 where the aqueous medium is selected from the group consisting of water; buffered water; phosphate buffered water; phosphate buffered saline; citrate buffered water; acetate buffered water; water buffered with pharmaceutically acceptable pH controlling agents; water containing salts; water containing sodium chloride; water containing pharmaceutically acceptable salts; water containing soluble agents for lyoprotection; water containing soluble agents for cryoprotection; water containing dextrose; water containing mannitol; water containing trehalose; water containing sucrose; water containing sorbitol; water containing pharmaceutically acceptable lyoprotectants; water containing pharmaceutically acceptable cryoprotectants; water containing polyhydroxy-containing compounds; water containing sugars; water containing polyols, and mixtures thereof.
14. The compositions of claims 1 and 2 where the aqueous medium is selected from the group consisting of a biological fluid; blood; plasma; saliva; urine; a protein-containing solution; an aqueous suspension of a protein; lymph fluid; semen; vaginal fluid; lachrymal fluid; nasal fluid; synovial fluid; cerebral fluid; cerebrospinal fluid; amniotic fluid; pancreatic fluid; pulmonary fluid; ascites fluid; fluid from a cyst; gastric fluid; intestinal fluid; a fluid removed from a patient; a diluted biological fluid; a concentrated biological fluid; and a mixture of biological fluids from one or more patients.

SURFACE MODIFIED PARTICULATE COMPOSITIONS OF BIOLOGICALLY ACTIVE SUBSTANCES

ABSTRACT

This invention disclosure relates to compositions for the delivery of stable surface modified sub-micron and micron sized particles of water-insoluble biologically active substances from a non-aqueous medium that self-disperses on exposure to an aqueous environment.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
220

Figure 1




Figure 2


Process Flow Diagram

Figure 3


Process Flow Diagram

Figure 4


Process Flow Diagram

Figure 5

Process Flow Diagram